

## Family 84. AIZOACEAE

### Revision:

Backer, CA. 1951. *Aizoaceae*. Flora Malesiana series 1, 4: 271-275

### 1. *Sesuvium* Linnaeus

***Sesuvium portulacastrum*** (L.) L., Syst. ed. 10 (1759) 1058; --Merr., Fl. Manila (1912) 88; EPFP 2 (1923) 136; --Backer, Fl. Males ser. 1, 4 (1951) 272, Fig. 2.

Pantropic. Throughout Malesia but not yet reported for Borneo. In saline, clayey or sandy, humid or muddy sites near the sea and along tidal creeks above 0-1m above sea level, often in dense patches.

### 2. *Trianthema* Linnaeus

***Trianthema portulacastrum*** L., Sp. Pl. (1753) 223; --Merr., Fl. Manila (1912) 198; EPFP 2 (1923) 136; --Backer, Fl. Males ser. 1, 4 (1951) 274.

Pantropic. In Malesia: Peninsular Malaysia, Java, Philippines, Madura, Lesser Sunda Isls (Sumbawa, Flores, Timor), Moluccas (Ternate), S New Guinea.

***Trianthema triquetra*** Rottl. ex Willd., Neue Schr. Naturfr. Berlin 4 (1803) 181; --Backer, Fl. Males. ser. 1, 4 (1951) 273.

Tropical Africa and Asia to Victoria, in Malesia: NE Java and Lesser Sunda Isls (Bali); Philippines: MINDANAO (*fide* Steen., Blumea 12 (1964) 320.

### 3. *Tetragonia* Linnaeus

***Tetragonia tetragonoides*** (Pallas) Kuntze, Rev. Gen. (1891) 264; --Backer, Fl. Males ser. 1, 4 (1951) 275.

Wild in the coastal regions of Australia, Tasmania, New Zealand, the Pacific islands, and Japan. Cultivated mostly 1000-1700m in Malesia. New Zealand spinach.