

Family 130. OXALIDACEAE

Revision:

Veldkamp, JF. 1971. *Oxalidaceae*. Flora Malesiana series 1, 7:

Averrhoa Linnaeus

Averrhoa bilimbi L., Sp. Pl. (1753) 428; --Merr., Fl. Manila (1912) 266; EPFP 2 (1923) 324;

Averrhoa carambola L., Sp. Pl. (1753) 428; --Merr., Fl. Manila (1912) 266; EPFP 2 (1923) 324;

Biophytum A de Candolle

Biophytum fruticosum Blume, Bijdr. (1825) 242;

var. fruticosum

NE India (Silhet), SW China, Hainan, Indochina (Mekong) and Malesia: E Sulawesi, S C Timor and the Philippines: BOHOL, SULU and BENGAL Is.

Biophytum microphyllum Veldk., Fl. Males. ser. 1, 7 (1971) 164.

SW & C Sulawesi, Flores and Philippines, MINDORO: Mindoro Oriental (Mt Yagaw).

Biophytum petersianum Klotzsch in Peters, Reise Mossamb. Bot. 1 (1862) 81, t. 15; --Veldk., Fl. Males. ser. 1, 7 (1971) 161.

Biophytum apodiscias (Turcz.) Edgew. & Hook.f. in Hook.f., Fl. Brit. Ind. 1 (1874) 437; --Merr., EPFP 2 (1923) 323.

Biophytum sensitivum (L.) DC, Prodr. 1 (1824) 690; --Merr., Fl. Manila (1912) 266; EPFP 2 (1923) 324.

Oxalis Linnaeus

Oxalis acetosella L., Sp. Pl. (1753) 433;

ssp. griffithii (Edgew. & JD Hook.) Hara, J. Jap. Bot. 30 (1955) 22; --Veldk., Fl. Males. ser. 1, 7 (1971) 157.

Khasya, Sikkim, Bhutan through China to Japan and Taiwan. In Malesia so far only known from Philippines, LUZON: Benguet prov., Mt Pulog, 2 coll.

Oxalis corniculata L., Sp. Pl. (1753) 435; --Veldk., Fl. Males. ser. 1, 7 (1971) 155.

Oxalis repens Thunb., Diss. Oxal. (1781) 16; --Merr., Fl. Manila (1912) 265; EPFP 2 (1923) 323.

Oxalis corymbosa DC, Prodr. 1 (1824) 696; --Veldk., Fl. Males. ser. 1, 7 (1971) 159.

Oxalis martiana Zucc., Denkschr. Akad. Muench. 9 (1823-24) 144; --Merr., Fl. Manila (1912) 265; EPFP 2 (1923) 323.

Doubtful species:

Sarcotheca philippica (Fern.-Villar) Hallier f., Meded. Rijksherb. (1911) 2;