

## Family 37. STEMONACEAE

### Revision:

Duyfjes, BEE. 1993. Stemonaceae. Flora Malesiana series 1, 11: 399-409.

### 1. *Stemona* Loureiro

**1. *Stemona lucida*** (R. Br.) Duyfjes, Blumea 36 (1991) 243; Fl. Males. ser. 1, 11 (1993) 404. Type from

Queensland.

*Stemona philippinensis* Merr., Govt. Lab. Publ. (Philipp.) 6 (1904) 16; EPFP 1 (1922) 201. --Type: Merrill 3061 (K,

holotype). Masbate.

Australia (N Queensland), Papua New Guinea, and the Philippines: MASBATE, TICA0.

Rather dry places in monsoon areas, in scrub, open forest, dry gallery forest, an on roadsides, alt. 0-200m.

**2. *Stemona tuberosa*** Lour., Fl. Cochinch. (1790) 404; --Merr., EPFP 1 (1922) 202; --Buysen, Fl. Males. ser. 1,

11 (1993) 405, f. 1-2. --Type: No specimen found, the illustration in Rumph., Herb. Amb. 5 (1747) 365 "Ubi

Gorita nigrum", t. 129, serving as lectotype.

#### **var. *tuberosa***

SE Continental Asia, Taiwan, Hainan, Philippines, Flores, Ambon.

**var. *ternatensis*** (JJ Sm.) Duyfjes, Blumea 36 (1991) 243; Fl. Males. ser. 1, 11 (1993) 405, f. 1-2. --Type from Ternate.